



Original Research

Effect of Feeding Rice Distillers Dried Grains with Solubles and Rice Gluten Meal with or without Enzyme Supplementation on the Haematology and Serum Biochemistry of Broiler Chickens

Om Prakash Dinani*, Pramod Kumar Tyagi, Nasir Akbar Mir, Asit Baran Mandal, Praveen Kumar Tyagi, Sita Prasad Tiwari¹ and Anil Kumar Giri

ICAR-Central Avian Research Institute Izatnagar, Bareilly-243122, Uttar Pradesh, INDIA

¹College of Veterinary Science and Animal Husbandry, Anjora, Durg-491001, Chhattisgarh, INDIA

*Corresponding author: dr_dinani@rediffmail.com

Rec. Date:	Jul 17, 2018 03:38
Accept Date:	Oct 08, 2018 17:39
DOI	10.5455/ijlr.20180717033823

Abstract

A biological experiment of 42 days duration was undertaken in which day old 240 chicks were divided into six dietary treatments as per 2x3 factorial design having 5 replicates per treatment with 8 birds in each to evaluate the effect of feeding rice distillers dried grains with solubles (rDDGS) and rice gluten meal (RGM) combination as soybean replacement along with exogenous enzyme supplementation on the haematology and serum biochemistry of broiler chicken at 21 days of age. The results revealed that no significant ($P>0.05$) differences were observed in blood profile between control and other different dietary treatments. Serum glucose, total protein and albumen levels decreased significantly ($P<0.01$) in rDDGS and RGM combinations as compared to control while other serum enzymes and serological parameters showed insignificant ($P>0.05$) effect due to feeding of rDDGS and RGM in combination up to the levels of 12.5% and 15%, respectively with or without enzyme supplementation.

Key words: Broiler, DDGS, Hematology, Protease, Rice Gluten Meal, Serum Biochemistry

How to cite: Dinani, O., Tyagi, P., Mir, N., Mandal, A., Tyagi, P., Tiwari, S., & Giri, A. (2019). Effect of Feeding Rice Distillers Dried Grains with Solubles and Rice Gluten Meal with or without Enzyme Supplementation on the Haematology and Serum Biochemistry of Broiler Chickens. International Journal of Livestock Research, 9(1), 254-261. doi: 10.5455/ijlr.20180717033823

Introduction

The diet and plane of nutrition are prominent factors that influence the blood picture and serum biochemistry of animals, which are sensitive indicators of the state of health and reflect the intensity of metabolic processes taking place in the animals (Vazquez-Anon *et al.*, 2008; Andrews *et al.*, 2006; Roy *et al.*, 2013). Similarly, it has been reported that the serum biochemical parameters are the good indicators of the physiological, nutritional, and pathological status of animals/birds and can be correlated to identify the





impact of nutritional factors and additives supplied in the diet (Toghyani *et al.*, 2010). Thus, the dietary manipulations in broiler chicken are expected to reveal differential effects on the blood picture and serum biochemistry of broiler chicken. In this regard, the rice distillers dried grains with solubles (rDDGS) and rice gluten meal (RGM) are two potential economic feed ingredients of poultry ration produced by rice processing industries. India is one of the largest producers of rice in world producing approximately 105 million tons of rice in 2015-16 (Govt. of India, 2016). The rising cost of broiler chicken ration can be overcome by the use of 'rice DDGS and rice gluten meal', protein rich alternative feed ingredients to replace costly soybean meal. An increase in ethanol production over the last 5-10 years, due to increasing prices of conventional oil and limited underground reserves, has led to an increased supply of DDGS that may be available as livestock feed (Noll *et al.*, 2007). On the other hand, the RGM, a by-product of wet-milling of rice, is available in appreciable amounts at lower cost compared to soybean. RGM contains 3152 kcal ME/kg, 46.45% crude protein, 3.4% ether extract and a favorable amino acid profile with relatively higher abundance of methionine (Kumar *et al.*, 2016). It has been also designated as a source of rumen undegradable protein with the highest known metabolizable protein value among plant proteins (Heuze *et al.*, 2015). Further, the exogenous enzyme supplementations in poultry diets, nutritionally, economically and environmentally justified, increase nutrient digestibility, reduce water content and viscosity of the excreta, and accelerate the rate of passage of digesta through the gastrointestinal tract (Sarmiento-Franco *et al.*, 2003; Lazaro *et al.*, 2004). In this study, it is hypothesised that the use of rDDGS and RGM in combination up to 12.5 and 15% levels respectively along with enzyme supplementation will not have any significant effect on the blood parameters and serum biochemistry of broiler chicken at 21 days of age. Thus, to test this hypothesis the objective of this study was to evaluate the effects of feeding different combinations of rDDGS and RGM levels along with enzyme supplementation on the haematology and serum biochemical parameters of broiler chicken.

Materials and Methods

Birds, Experimental Diets and Design

A total of 240 CARIBRO Vishal broiler chicks of same hatch and uniform weight were procured from the institutional hatchery and housed in specially designed battery brooder cages with standard watering and feeding facilities. A biological experiment of 42 days duration was undertaken in which day old 240 chicks were divided into six dietary treatments as per 2x3 factorial design having 5 replicates per treatment with 8 birds in each. The broiler chicken ration were formulated by using rDDGS and RGM as replacement of soybean meal in the basal diets along with enzyme supplementation (ICAR, 2013). The levels of rDDGS and RGM and the suitable enzymes were standardised in a preliminary trail. The rDDGS levels of 10% & 12.5%, and RGM levels of 12.5% & 15% along with either multienzyme or protease enzyme were selected



to formulate the experimental diets as the pre-starter, starter and finisher diets. The feed ingredients and the nutrient composition of diets have been given in Table 1. A 2x3 factorial design resulted in six experimental diets viz T1 (no rDDGS/RGM/enzyme), T2 (no rDDGS/RGM, with multienzymes), T3 (12.5% rDDGS, 15% RGM, no enzyme), T4 (12.5% rDDGS, 15% RGM, with protease enzyme), T5 (10% rDDGS, 12.5% RGM, no enzyme), and T6 (10% rDDGS, 12.5% RGM, with protease enzyme). Each treatment was allocated 5 replicates of chicks, with 8 birds in each. The feeding trial was conducted for six weeks and the feed as well as drinking water were provided *ad libitum* to the birds during the entire experimental period.

Table 1: Ingredient and nutrient composition of broiler chicken diets

Ingredient	Pre-starter diet (0-14days)						Starter diet (14-28days)						Finisher diet (28-42days)					
(%)	T1	T2	T3	T4	T5	T6	T1	T2	T3	T4	T5	T6	T1	T2	T3	T4	T5	T6
Maize	54.42	54.42	58.58	58.58	57.78	57.78	55.63	55.63	60.97	60.97	59.81	59.81	62	62	66.77	66.77	65.61	65.61
SBM	38.4	38.4	8.8	8.8	14.3	14.3	37.1	37.1	7.4	7.4	13	13	31.3	31.3	2	2	7.5	7.5
DORB	0	0	0.7	0.7	0.9	0.9	0	0	12.5	12.5	10	10	3.22	3.22	0	0	0.7	0.7
DDGS	0	0	12.5	12.5	10	10	0	0	15	15	12.5	12.5	0	0	12.5	12.5	10	10
RGM	0	0	15	15	12.5	12.5	3.5	3.5	0.2	0.2	0.8	0.8	0	0	15	15	12.5	12.5
Oil	3	3	0	0	0.4	0.4	1.35	1.35	1.15	1.15	1.23	1.23	0.5	0.5	0.5	0.5	0.5	0.5
LSP	1.4	1.4	1.3	1.3	1.1	1.1	1.55	1.55	1.7	1.7	1.67	1.67	0.7	0.7	0.4	0.4	0.33	0.33
DCP	1.82	1.82	2	2	2	2	0	0	0.32	0.32	0.22	0.22	1.45	1.45	1.7	1.7	1.64	1.64
Lysine	0	0	0.35	0.35	0.25	0.25	0.1	0.1	0	0	0	0	0	0	0.36	0.36	0.3	0.3
Methionine	0.2	0.2	0	0	0	0	0.765	0.765	0.765	0.765	0.765	0.765	0.06	0.06	0	0	0	0
Constant*	0.765	0.765	0.765	0.765	0.765	0.765	55.63	55.63	60.97	60.97	59.81	59.81	0.77	0.77	0.77	0.77	0.77	0.77
Enzyme	--	M	--	P	--	P	--	M	--	P	--	P	--	M	--	P	--	P
Nutrient Composition																		
CP (%)	21.99	21.99	22.02	22.02	22.02	22.02	21.52	21.52	21.5	21.5	21.49	21.49	19.51	19.51	19.52	19.52	19.49	19.49
ME (kcal/kg)	2998	2998	3002	3002	2998	2998	3050	3050	3051	3051	3050	3050	3100	3100	3099	3099	3104	3104
Ca (%)	1.03	1.03	1.08	1.08	1	1	0.95	0.95	0.95	0.95	0.96	0.96	0.86	0.86	0.85	0.85	0.86	0.86
Available P (%)	0.45	0.45	0.45	0.45	0.46	0.46	0.41	0.41	0.4	0.4	0.4	0.4	0.38	0.38	0.39	0.39	0.39	0.39
Lysine (%)	1.19	1.19	1.19	1.19	1.2	1.2	1.38	1.38	1.12	1.12	1.13	1.13	1.2	1.2	1	1	1.04	1.04
Methionine (%)	0.52	0.52	0.53	0.53	0.51	0.51	0.48	0.48	0.53	0.53	0.5	0.5	0.41	0.41	0.5	0.5	0.48	0.48
Threonine (%)	0.8	0.8	0.81	0.81	0.82	0.82	0.79	0.79	0.81	0.81	0.8	0.8	0.86	0.86	0.79	0.79	0.81	0.81
Cost (Rs./kg)	28.52	28.93	23.02	23.63	23.68	24.29	28.03	28.43	22.88	23.48	23.65	24.25	26.72	26.72	22.03	22.03	22.93	22.93

SBM: Soybean meal, DORB: Deoiled rice bran, DDGS: Dried distillers grains with solubles, RGM: Rice gluten meal, LSP: Limestone powder, DCP: Di-calcium phosphate, CP: Crude protein, ME: Metabolizable energy; *Constant (0.4% salt, 0.1% trace mineral premix, 0.15% vitamin premix, 0.015% vit. B complex, 0.05% choline chloride and 0.05% Toxin binder); 1: Trace mineral premix supplied (mg/kg diet): Mg 300; Mn 55; I 0.4; Fe 56; Zn 30; Cu 4.; 2: Vitamin premix supplied (per kg diet): Vitamin A 8250 IU; Vitamin D3 1200 IU; Vitamin K 1mg; Vitamin E 40 IU.; 3: B complex: Vitamin B1 2 mg; Vitamin B2 4 mg; Vitamin B12 10 µg; niacin 60 mg; pantothenic acid 10 mg; choline 500 mg.

Sampling and Measurement

All the procedures carried out on the birds were approved by the Institutional Animal Ethics Committee of Indian Veterinary Research Institute, Izatnagar (452/01/ab/CPCSEA). For the analysis of haematological parameters 2 ml blood samples were collected from 8 birds (4 males and 4 females) per treatment at random in sterile heparin vials at 21 days of age of birds. All the blood samples were analyzed by automatic Abacus



junior vet 5 haemato analyzer. The parameters studied were total leucocyte count (TLC), differential leucocyte count (DLC), Hb (%), packed cell volume (PCV), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), RBC distribution width (RDWc), platelet count, mean platelet volume (MPV) and platelet distribution width (PDWc). For evaluation of serum biochemistry of broiler chicken at 21 days of age, the blood samples were collected in the same manner as for the haematological analysis but in vials without any anticoagulant. The various serological parameters studied were serum glucose (Trinder, 1969), total protein (Doumas *et al.*, 1971), albumin (Gustafsson, 1978), globulin, albumin: globulin ratio (A:G), cholesterol (Wybenga and Pileggi, 1970), triglyceride (Fossati and Lorenzo, 1982), SGOT (Reitman and Frankel, 1957), SGPT (Reitman and Frankel, 1957) and ALP (Kind and King, 1954).

The data collected were analysed by two way ANOVA using GLM procedure to present the results as means and standard errors (Snedecor and Cochran, 1989) by using statistical package for social sciences (SPSS) 16.0 version and the comparison of significant mean differences was done by Duncan's multiple range test (Duncan, 1955) with significance level set at $P < 0.05$.

Results and Discussion

The results of feeding different level of rDDGS and RGM combinations with or without enzymes on haematological parameters of the broiler chicken of 21 days of age are given in Table 2. The blood profile was studied in terms of total erythrocyte count (TEC), total leukocyte count (TLC), differential leukocyte count (DLC), platelet count, Hb %, PCV, MCV, MCH, MCHC, MPV, heterophils and leukocyte ratio, RBC distribution width (RDWc) and platelet distribution width (PDWc). The results revealed that no significant ($P > 0.05$) differences were observed in blood profile between control and other different dietary treatments having different levels of rDDGS and RGM combinations with or without enzymes and their interactions. All the blood parameters studied were well within the normal physiological range. In line with the results of the present study, Ghazalah *et al.* (2011) and Youssef *et al.* (2013) reported no significant effect of DDGS inclusion up to 15% level. Similarly, Wani *et al.* (2017) reported no significant ($P > 0.05$) difference in Hb and PCV of broiler chicken by incorporating RGM up to 17.5% level with or without protease supplementation. However, in contrast to the results of the present study Gupta *et al.* (2017) reported inclusion of rDDGS level above 5% level significantly ($P < 0.01$) enhance the PCV and Hb values of broiler chicken compared to control diet fed birds.

Table 2: Effect of feeding different level of rDDGS and RGM combinations haematological parameters

rDDGS	RGM	Enzyme	TLC	Hetero.	Lymph	Mono	Hetero. %	Lymph %	Mono %	H:L	TEC	Hb	PCV	MCV	MCH	MCHC	RDWc	PLT	PLT%	MPV	PDWc
0.00%	0.00%	-	15.8	3.7	9.9	0.9	24	62.6	5.7	0.38	2.8	11.4	25.8	91	40.5	44.5	13.5	28.2	0.28	7.9	33.5
0.00%	0.00%	M	15.6	3.5	9.7	0.8	23.1	62.5	5.6	0.37	2.8	11.3	25.3	89.7	40.2	44.5	13.2	26	0.19	8.2	34
12.50%	15.00%	-	16.1	3.7	9.9	0.9	23	61.9	5.9	0.37	2.7	11.1	24.7	89.2	40	44.7	14.5	27	0.27	7.3	31.5
12.50%	15.00%	P	14.7	3.3	9	0.8	22.9	61.5	5.8	0.38	2.8	11.1	25.5	89.2	40	44	13.2	27.2	0.27	7.9	31.5
10.00%	12.50%	-	14.8	3.6	8.6	0.8	24.1	58	5.3	0.42	2.8	11	25.6	89.5	39.8	43.2	13	28.5	0.29	7.6	33.7
10.00%	12.50%	P	16.7	3.8	9.5	1	23.1	63	5.9	0.37	2.8	11.9	26.5	92.5	41.7	45	13.5	28	0.28	7.6	32.5
Pooled SEM			0.31	0.07	0.23	0.03	0.34	0.59	0.16	0.01	0.02	0.13	0.25	0.62	0.29	0.32	0.19	0.61	0.01	0.11	0.34
Main Effects (rDDGS/RGM combination)																					
0.00%	0.00%		15.8	3.7	9.9	0.8	23.7	62.8	5.4	0.38	2.8	11.3	25.6	91	40.5	44.4	13.4	27.8	0.24	8	33.7
12.50%	15.00%		15.3	3.5	9.4	0.8	22.8	61.4	6	0.37	2.8	11.1	25.1	88.8	39.8	44.4	13.7	26.5	0.27	7.6	31.7
10.00%	12.50%		15.8	3.7	9	0.9	23.6	60.5	5.6	0.39	2.8	11.5	26	91	40.7	44.1	13.2	28.2	0.28	7.6	33.1
Enzyme Supplementation																					
Without enzyme			15.6	3.6	9.5	0.8	23.7	60.9	5.6	0.3	2.8	11.2	25.4	89.9	40.1	44.1	13.6	27.9	0.28	7.6	32.9
With enzyme			15.7	3.6	9.4	0.8	23	62.2	5.8	0.3	2.8	11.4	25.8	90.5	40.6	44.5	13.3	27	0.25	7.9	32.6
Significance																					
rDDGS/RGM combination			NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Enzyme supplementation			NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Interaction			NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Values bearing different superscripts within the column differ significantly; NS: Non-significant ($P>0.05$); Units: WBC ($\times 10^4$ ul), RBC ($\times 10^6$ ul), Hb (g/dl), MCV (fl), MCH(pg), MCHC(g/dl) and PLT($\times 10^3$ ul)



The results pertaining to feeding different level of rDDGS and RGM combinations on serological parameters on 21 days are presented in Table 3. The results revealed that no significant ($P>0.05$) difference was found in serum globulin (G), albumen(A): globulin (G) ratio, serum enzymes SGOT, SGPT and ALP activity between different dietary treatments and control as result of feeding 0% rDDGS +0% RGM (DR1), 12.5% rDDGS +15% RGM (DR2) and 10 % rDDGS +12.5 % RGM (DR3) combination levels. Serum glucose, total protein and albumen levels decreased significantly ($P<0.01$) in DR2 and DR3 as compared to DR1 level. However, serum glucose, total protein and albumen levels did not show any significant ($P>0.05$) difference from DR2 as compared to DR3. Serum triglyceride levels were decreased significantly ($P<0.01$) in DR2 level as compared to DR1 and DR3 levels. Serum triglyceride level did not show any significant ($P>0.05$) difference from DR3 as compared to DR1 level. Effect of with or without enzymes supplementation on feeding different level of rDDGS and RGM combinations did not show any significant ($P>0.05$) difference in serological parameters except serum glucose. Serum glucose level was increased significantly ($P<0.01$) in enzymes supplemented groups as compared to without enzymes groups. Interaction of rDDGS and RGM combinations with or without enzymes did not show any significant ($P>0.05$) difference on serological parameters on 21 days of broiler chickens. Our results were in agreement with Choi *et al.* (2014), Metwally and Farhat (2015), Kumar *et al.* (2016) and Wani *et al.* (2017). Choi *et al.* (2014) reported dietary inclusion of rice DDGS up to 25% did not affect the plasma content of total protein, glucose, cholesterol and triglyceride in juvenile red sea bream (*Pagrus major*) fish. Metwally and Farhat (2015) reported no significant ($P>0.05$) difference in serum biochemical parameters (serum lipid profile, glucose, total protein, albumin and globulin) up to addition of 12.5% RGM in diet of broiler chicken. Kumar *et al.* (2016) also reported no significant ($P>0.05$) difference in the serological variables (glucose, blood urea nitrogen, plasma proteins and non-esterified fatty acids) on addition of RGM up to 21% level in the diet of growing dairy calves. Wani (2017) and Wani *et al.* (2017) reported no significant ($P>0.05$) difference on feeding different levels of RGM up to 20% level with or without protease enzyme supplementation. Moreover, serum biochemical parameters are indicators of the physiological, nutritional and pathological status of birds and can be correlated to identify the impact of nutritional factors and additives supplied in the diet.

Our results were in disagreement with Hack *et al.* (2015) and Gupta *et al.* (2017). Hack *et al.* (2015) reported that increasing corn DDGS level significantly ($P<0.01$) increased serum triglycerides, cholesterol and LDL for hens fed diet contained 22% DDGS in the diet. Gupta *et al.* (2017) reported 10% rice DDGS had significantly ($P<0.01$) higher effect on serum albumin, total protein, A:G ratio, glucose value and significantly ($P<0.01$) lowering effect on serum lipid profile (triglycerides, cholesterol, LDL and VLDL). However, no such study on rDDGS and RGM combination feeding were done in poultry on blood biochemical parameters till now.

Table 3: Effect of feeding different level of rDDGS and RGM combinations on serological parameters

rDDGS	RGM	Enzyme	Glucose mg/dl	Protein (g/dl)	Albumin g/dl	Globulin g/dl	A:G	Cholesterol mg/dl	Triglyceride mg/dl	SGOT U/ml	SGPT U/ml	ALP U/ml
0.00%	0.00%	-	183	4.4	1.9	2.3	0.68	105	97	165	12	50
0.00%	0.00%	M	191	4.6	2	2.5	0.66	104	90	164	13	57
12.50%	15.00%	-	171	4.2	1.6	2.5	0.7	103	81	167	15	63
12.50%	15.00%	P	186	4.1	1.6	2.5	0.67	99	84	165	13	55
10.00%	12.50%	-	174	3.9	1.7	2.6	0.73	105	91	164	12	52
10.00%	12.50%	P	184	4	1.7	2.7	0.69	105	89	166	13	54
Pooled SEM			1.881	0.037	1.44	0.05	0.02	0.03	0.01	0.88	1.49	0.54
Main Effects (rDDGS/RGM combination)												
0.00%	0.00%		187 ^b	4.5 ^b	1.9 ^b	2.4	0.67	105	94 ^b	164	13	53
12.50%	15.00%		178 ^a	4.2 ^a	1.6 ^a	2.5	0.69	101	83 ^a	166	14	59
10.00%	12.50%		179 ^a	4.0 ^a	1.7 ^a	2.6	0.71	105	90 ^b	165	12	53
Enzyme Supplementation												
Without enzyme			176 ^a	4.2	1.7	2.5	0.7	104	90	165	13	55
With enzyme			187 ^b	4.2	1.7	2.5	0.67	103	88	165	13	55
Significance												
rDDGS/RGM combination			P<0.01	P<0.01	P<0.01	NS	NS	NS	P<0.01	NS	NS	NS
Enzyme supplementation			P<0.01	NS	NS	NS	NS	NS	NS	NS	NS	NS
Interaction			NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Values bearing different superscripts within the column differ significantly; A:G ratio: Albumin: Globulin ratio, NS: Non-significant ($P>0.05$); M: Multienzyme, P: Protease

Conclusion

The study concludes that the feeding of rDDGS and RGM in combination up to the levels of 12.5% and 15%, respectively with or without enzyme supplementation do not have any adverse effects on the haematological parameters but adversely affecting serum biochemical parameters in terms of serum glucose, total protein and albumen levels of broiler chicken in 21 days.

Acknowledgement

Authors are highly thankful to ICAR-Central Avian Research Institute, Izatnagar, Bareilly, U.P. for providing all necessary facilities and inputs.

References

- Andrews J, Vazquez-Anon M, and Bowman G. 2006. Fat stability and preservation of fatty acids with AGRADO antioxidant in feed ingredients used in ruminant rations. *Journal of Dairy Science*, 89:60-69.
- Choi J, Rahman MM, and Lee SM. 2014. Rice Distillers Dried Grain is a Promising Ingredient as a Partial Replacement of Plant Origin Sources in the Diet for Juvenile Red Seabream (*Pagrus major*). *Asian-Australian Journal of Animal Science*, 27(12): 1736–1743.
- Doumas BT, Watson WA, and Biggs HG. 1971. Albumin standards and the measurement of serum albumin with bromocresol green. *Clinica Chimica Acta*, 31: 87-96.
- Duncan DB. 1955. Multiple range and F tests. *Biometrics*, 11: 1-42.
- Fossati P and Lorenzo P. 1982. Serum triglycerides determined calorimetrically with an enzyme that produces hydrogen peroxide. *Clinical Chemistry*, 28: 2077-80.
- Ghazalah AA, Abd-Elsamee MO, and AL-Arami AA. 2011. Use of distillers dried grains with solubles (DDGS) as replacement for yellow corn in laying hen diets. *Egyptian Poultry Science*, 31: 191-202.



7. Govt. of India, Ministry of Agriculture & Farmers Welfare. 2016. Area, production and yield of principal crops. In: Agricultural Statistics at a glance 2016. Govt. of India Controller of Publication. pp. 87-88.
8. Gupta, S.L., Tyagi, Pramod K., Tyagi, Praveen K., Mandal, A.B., Dinani, O.P. and Rokade, J.J. 2017. Feeding effect of rice based dry distillers grains with soluble on hemato-biochemical and egg sensory attributes during 45th to 54th week of laying. *International Journal of Pure Applied Biosciences*, 5(6): 1521-1527.
9. Gustafsson JE. 1978. Automated serum albumin determination by use of the immediate reaction with bromocresol green reagent. *Clinical Chemistry*, 24(2): 369-373.
10. Hack, MA, El-Hindawy, MM, Attia, AI and Khalid, M 2015. Effects of feeding dried distillers grains with solubles with or without enzyme or vitamin E supplementation on productive performance of hisex brown laying hens. *Zagazig Journal of Agricultural Research*, 42(1):71-79.
11. Heuze V, Tran G, Sauvant D, Renaudeau D, Lessire M, and Lebas F. 2015. Corn gluten feed. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <https://www.feedipedia.org/node/714>
12. ICAR. 2013. Nutrient requirements of Animals-Poultry (ICAR-NIANP). 13-16.
13. Kind PRN and King EJ. 1954. Estimation of Plasma Phosphatase by Determination of Hydrolysed Phenol with Amino-antipyrine. *Journal of Clinical Pathology*, 7(4): 322-326.
14. Kumar R, Thakur SS, and Mahesh MS. 2016. Rice gluten meal as an alternative by product feed for growing dairy calves. *Tropical Animal Health Production*, 48(3):619-624.
15. Lazaro R, Latorre MA, Medel P, Gracia M, and Mateos GG. 2004. Feeding regimen and enzyme supplementation to rye based diets for broilers. *Poultry Science*, 83:152-160.
16. Metwally A and Farahat M. 2015. Nutritive value and feeding of rice gluten meal in broiler chickens. *Research Opinion in Animal and Veterinary Sciences*, 5(11): 443-451.
17. Noll SE, Parsons CM, and Dozier III WA. 2007. Formulating poultry diets with DDGS-How far can we go. In: Proceedings 5th Mid Atlantic Nutrition Conference. Zimmerman, NG. ed., University of Maryland, College Park, USA, 91-99.
18. Reitman S and Frankel S. 1957. A colorimetric test for determination of serum glutaminoxaloacetic and glutamic pyruvic transaminase. *American Journal of Clinical Pathology*, 28: 56-63.
19. Roy A, Mandal G, and Patra A. 2013. Evaluating the performance, carcass traits and conjugated linoleic acid content in muscle and adipose tissues of black Bengal goats fed soybean oil and sunflower oil. *Animal Feed Science and Technology*, 185:43-52.
20. Sarmiento-Franco L, McNab JM, Pearson A, and Belmar-Casso R. 2003. The effects of chaya leaf meal and of exogenous enzymes on amino acid digestibility in broilers. *British Poultry Science*, 44: 458-63.
21. Snedecor GW and Cochran WG. 1989. Statistical Methods. 7th edn. Oxford and IBH.
22. Toghyani M, Tohidi M, Gheisari A.A, Tabeidian SA 2010. Performance, immunity, serum biochemical and hematological parameters in broiler chicks fed dietary thyme as alternative for an antibiotic growth promoter. *African Journal of Biotechnology*, 9(40): 6819-6825.
23. Trinder P. 1969. Enzymatic methods for glucose determination. *Annals of Clinical Biochemistry*, 6: 24-26.
24. Vazquez-Anon M, Nocek J, Bowman G, Hampton T, Atwell C, Vazquez P, and Jenkins T. 2008. Effects of feeding a dietary antioxidant in diets with oxidized fat on lactation performance and antioxidant status of the cow. *Journal of Dairy Science*, 91: 3165-3172.
25. Wani MA, Tyagi Pramod K, Tyagi Praveen K, Sheikh SA, Dinani OP, Hazarika R, Bhanja SK, and Mandal AB. 2017. Effect of rice gluten meal as protein source in the diet of broiler chicken on immunity, gut microbial count, haematology and serum biochemical parameters. *Indian Journal of Poultry Science*, 52(3): 277-282.
26. Wybenga DR and Pileggi VJ. 1970. Estimation of Cholesterol. *Clinical Chemistry*, 16: 980.
27. Youssef AW, El-Azeem NAA, El-Daly EF, and Monairy MM. 2013. The Impact of Feeding Graded Levels of Distillers Dried Grains with Solubles (DDGS) on Broiler Performance, Haematological and Histological Parameters, *Asian Journal of Poultry Science*, 7 (2): 41-54.

